



Pacific Country Report

Sea Level & Climate: Their Present State

Vanuatu

June 2002



This project is sponsored by the Australian Agency for International Development (AusAID), managed by Australian Marine Science and Technology Ltd (AMSAT), and supported by NTF Australia at the Flinders University of South Australia.

<u>Disclaimer</u> The views expressed in this publication are those of the authors and not necessarily those of the Australian Agency for International Development (AusAID)

PACIFIC COUNTRY REPORT ON SEA LEVEL & CLIMATE: THEIR PRESENT STATE



VANUATU

June 2002

Executive Summary

- A SEAFRAME gauge was installed in Port Vila, Vanuatu, in January 1993. It records sea level, air and water temperature, atmospheric pressure, wind speed and direction. It is one of an array designed to monitor changes in sea level and climate in the Pacific.
- This report summarises the findings to date, and places them in a regional and historical context.
- The sea level trend to date is +10.5 mm/year (as compared to a global average of 1-2 mm/year) but the magnitude of the trend continues to vary widely from month to month as the data set grows. A nearby gauge, with longer records but less precision and datum control, shows a trend of +6.21 mm/year.
- Variations in monthly mean sea level are dominated by the 1997/1998 El Niño, with a moderate seasonal cycle.
- Variations in monthly mean air and water temperature are likewise affected by the 1997/1998 El Niño, with pronounced seasonal cycles.
- A number of destructive Tropical Cyclones have passed near Vanuatu since the SEAFRAME was installed, and two in particular, in 1993 and 2001, have come close enough to Port Vila to have been recorded as very low barometric pressures. Both caused considerable damage.
- Two tsunamis, both due to undersea earthquakes in Vanuatu, have caused loss of life and damage to property in Vanuatu in the period since installation.
- The tsunami caused by the Peru earthquake of June 2001, which registered strongly on many Pacific SEAFRAME gauges, measured about 30 cm peak-to-trough at Port Vila.



National Tidal Facility Australia

Telephone: 8201 7532 Facsimile: 8201 7523

(ISD +618)

Email: ntf@flinders.edu.au

WWW: http://www.ntf.flinders.edu.au

Postal Address: GPO Box 2100, Adelaide 5001 Australia

Dear Pacific Island Government Representative

Welcome to the first **Pacific Country Report**, containing a summary of the sea level, climate, oceanography and extreme events for each of the twelve SEAFRAME monitoring sites, plus Palau and Niue. We intend to produce them to coincide with the Forum Meetings.

Your feedback is essential to ensure that improvements are made, that what is important to **you** is addressed and explained. Your feedback will help guide the frequency of publishing and distribution. We invite you to give us both positive and negative feedback (your comments will remain confidential) because what might be obvious to you might be overlooked by scientists.

You can tear out this page, jot notes on it, and mail or fax it to us at the address above. Or you can email comments to us. A few words is all we need.

1-Did you find it informative?
2-What significant information have we omitted?
3-Would you like to see additional emphasis on any topic? If so, what?
4-Would you like more explanation on any topic? If so, what?
5-Any other suggestions or constructive criticism?
Name (optional)

Thank you for your time!

Country

Introduction

As part of the AusAID-sponsored South Pacific Sea Level and Climate Monitoring Project ("Pacific Project") for the FORUM region, in response to concerns raised by its member countries over the potential impacts of an enhanced Greenhouse Effect on climate and sea levels in the South Pacific region, a **SEAFRAME** (**Sea** Level **F**ine **R**esolution **A**coustic **M**easuring **E**quipment) gauge was installed in Port Vila, Vanuatu, in January, 1993. Aside from a period following cyclone damage, the gauge has been returning high resolution, good scientific quality data since installation.

SEAFRAME gauges not only measure sea level by two independent means, but also a number of "ancillary" variables - air and water temperatures, wind speed, wind direction and atmospheric pressure. There is an associated programme of levelling to first order, to determine shifts in the vertical of the sea level sensors due to local land movement. Continuous GPS measurements are now also being made to determine the vertical movement of the land with respect to the International Terrestrial Reference Frame.

When change in sea level is measured with a tide gauge over a number of years one cannot be sure whether the sea is rising or the land is sinking. Tide gauges measure relative sea level change, i.e., the change in sea level relative to the tide gauge, which is connected to the land. To local people, the relative sea level change is of paramount importance. Vertical movement of the land can have a number of causes, e.g. island uplift, compaction of sediment or withdrawal of ground water. From the standpoint of global change it is imperative to establish absolute sea level change, i.e. sea level referenced to the centre of the Earth which is to say in the terrestrial reference frame. In order to accomplish this the vertical land movement and in particular the rate at which the land moves must be measured separately. This is the reason for the addition of CGPS near the tide gauges.

Regional Overview

Variations in sea level and atmosphere are inextricably linked. For example, to understand why the sea level at Tuvalu undergoes a much larger annual fluctuation than at Samoa, we must study the seasonal shifts of the trade winds. On the other hand, the climate of the Pacific Island region is entirely ocean-dependent. When the warm waters of the western equatorial Pacific flow east during El Niño, the rainfall, in a sense, goes with them, leaving the islands in the west in drought.

Compared to higher latitudes, air temperatures in the tropics vary little throughout the year. Of the SEAFRAME sites, the most extreme changes are naturally experienced by those furthest from the equator – the Cook Islands (at 21°S) recorded the lowest temperature, 13.1°C, in August 1998. The Cook Islands regularly fall to 16°C while Tonga (also at 21°S) regularly falls to 18°C in winter (July/August).

SEAFRAME location	Minimum recorded air temperature (°C)	Maximum recorded air temperature (°C)
Cook Islands	13.1	32.0
Tonga	16.0	31.4
Fiji (Lautoka)	16.6	33.4
Vanuatu	16.5	33.3
Samoa	18.7	32.1
Tuvalu	22.8	31.6
Kiribati	22.4	32.9
Nauru	22.4	32.4
Solomon Islands	20.1	34.5
Papua New Guinea	21.5	31.1
Marshall Islands	20.5	31.9

The most striking oceanic and climate fluctuations in the equatorial region are not the seasonal, but interannual changes associated with El Niño. These affect virtually every aspect of the system, including sea level, winds, precipitation, and air and water temperature. Referring to the plot below, we see that at most SEAFRAME sites, the lowest recorded sea levels appear during the 1997/1998 El Niño. The most dramatic effects were observed at the Marshall Islands, PNG, Nauru, Tuvalu and Kiribati, and along a band extending southeastward from PNG to Samoa. The latter band corresponds to a zone meteorologists call the "Sub-Tropical Convergence Zone" or STCZ. In the figure below, we see the effect of the 1997/1998 El Niño on all SEAFRAME stations.


Sea levels* at SEAFRAME sites

1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 Marshall Islands **Federated States of Micronesia** New site installed December 2001: 12 months of data needed for trend Papua New Guinea Solomon Islands Kiribati Nauru Tuvalu Samoa Vanuatu Fiji Tonga 0.2 0.2 Cook Islands 0.0 Dec Jun Dec Ju 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001

^{*} Plotted values are sea level "anomalies" (tides and trend removed from data).

Most Pacific Islanders are very aware that the sea level is controlled by many factors, some periodic (likes the tides), some brief but violent (like cyclones), and some prolonged (like El Niño), because of the direct effect the changes have upon their lives. The effects vary widely across the region. Along the Melanesian archipelago, from Manus Island to Vanuatu, tides are predominantly diurnal, or once daily, while elsewhere the tide tends to have two highs and two lows each day. Cyclones, which are fueled by heat stored in the upper ocean, tend to occur in the hottest month. They do not occur within 5° of the equator due to the weakness of the "Coriolis Force", a rather subtle effect of the earth's rotation. El Niño's impact on sea level is mostly felt along the STCZ, because of changes in the strength and position of the Trade Winds, which have a direct bearing on sea level, and along the equator, due to related changes in ocean currents. Outside these regions, sea levels are influenced by El Niño, but to a far lesser degree.


Mean Surface Water Temperature



Note the warm temperatures in the STCZ and just north of the equator.

The convergence of the Trade Winds along the STCZ has the effect of deepening the warm upper layer of the ocean, which affects the seasonal sea level. Tuvalu, which is in the heart of the STCZ, normally experiences higher-than-average sea levels early each year when this effect is at its peak. At Samoa, the convergence is weaker, and the seasonal variation of sea level is far less, despite the fact that the water temperature recorded by the gauge varies in a similar fashion. The interaction of wind, solar heating of the oceanic upper layer, and sea level, is quite complex and frequently leads to unexpected consequences.

The plot **Streamlines of Mean Surface Wind** shows how the region is dominated by easterly trade winds. In the Southern Hemisphere the Trades blow to the northwest and in the Northern Hemisphere they blow to the southwest. The streamlines converge, or crowd together, along the STCZ.



Much of the Melanesian subregion is also influenced by the Southeast Asian Monsoon. The strength and timing varies considerably, but at Manus Island (PNG), for example, the NW monsoon season (winds from the northwest) runs from November to March, while the SE monsoon brings wind (also known as the Southeast Trade Winds) from May to October. Unlike many monsoon-dominated areas, the rainfall at Manus Island is distributed evenly throughout the year (in normal years).

Mean Sea Level Trends and their Confidence Intervals

With the great diversity in climatic environments, vertical land movement and ocean variability, one might expect that that the sea level trends measured at different stations over the limited period for which tide gauge data has been collected may also vary. That this is indeed the case is demonstrated by the following table, which contains the relative sea level trends from all the regional stations for which at least 25 years of hourly data was available.


Location	Country	Years of	Trend	Standard
		data	(mm/year)	Deviation
				mm/year
Pago Pago	U S Trust	49.7	+1.43	1.5
Rarotonga	Cook Is	22.2	+3.80	3.7
Penrhyn	Cook Is	21.6	+0.89	3.4
Pohnpei	F S of Micronesia	26.9	+0.42	3.7
Kapingamarangi	F S of Micronesia	19.9	-1.04	4.7
Truk	F S of Micronesia	27.6	+1.79	3.3
Guam	U S Trust	50.1	+0.37	1.9
Yap	F S of Micronesia	30.9	-0.20	3.6
Suva	Fiji	24.8	+3.99	3.0
Christmas	Rep of Kiribati	40.3	-0.68	2.2
Kanton	Rep of Kiribati	45.0	+0.26	1.5
Fanning	Rep of Kiribati	16.8	+2.17	5.1
Tarawa	Rep of Kiribati	23.6	-2.24	3.6
Majuro	Rep of Marshall Is	30.8	+2.79	2.6
Enewetok	Rep of Marshall Is	24.5	+1.18	3.3
Kwajalein	Rep of Marshall Is	54.4	+1.13	1.3
Nauru	Rep of Nauru	24.2	-2.03	4.2
Malakal	Rep of Palau	30.1	+0.64	4.0
Honiara	Solomon Is	24.5	-2.21	4.8
Nuku'alofa	Tonga	9.4	+4.90	7.2
Funafuti	Tuvalu	21.6	+0.92	5.1
Port Vila	Vanuatu	11.3	+6.21	6.8

Mean trend: 1.11 mm/year (all data) Mean trend of data > 25 years: 0.8 mm/year Data from University of Hawaii as at June 2002

The following plot depicts the evolution of the short term sea level trends, at SEAFRAME stations, from one year after installation to the present. Please note that the trendlines have not yet stabilised.

Short Term Sea Level Trends (mm/year)


1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002



Dec Jun Dec Ju

The expected width of the 95% confidence interval (±1.96 times the standard error) as a function of data length based on the relationship for all National Oceanographic and Atmospheric Administration (NOAA) gauges with a data record of at least 25 years are shown in the figure below. A confidence interval or precision of 1 mm/year should be obtainable at most stations with 50-60 years of data on average, providing there is no acceleration in sea level change, vertical motion of the tide gauge, or abrupt shifts in trend due to tectonic events. In the figure, the 95% confidence intervals are plotted as a function of the year range of data, based on NOAA tide gauges with at least 25 years of record¹.

95% Confidence Intervals for Linear Mean Sea Level trends (mm/year)



This overview was intended to provide an introduction to the Pacific Islands regional climate, in particular those aspects that are related to sea level. This is an area of active research, and many elements, such as interdecadal oscillations, are only beginning to be appreciated. The individual country reports give greater detail on the variations experienced at the twelve SEAFRAME sites in the Pacific.

8

^{1.} Zervas, C. (2001) Sea Level Variations of the United States 1854-1999. NOAA, USA.

June-2002

Project findings to date - Vanuatu

Short-term sea level trend

A fundamental goal of the Project is to establish the rate of sea level change. It has been recognised since the beginning that this would require several decades of continuous, high quality data. However, in response to increasing requests from the region for information regarding the trends as they gradually emerge from the background "noise", combined with concern that less experienced users might attempt to fit a trend line to the data without properly accounting for processes such as seasonality that can bias the result, the preliminary findings are now being provided. These are given in the form of plots (see **Short Term Sea Level Trends** above) which show how the trend develops as more data becomes available. We caution against drawing conclusions prematurely.

As at June 2002, based on the short-term sea level rise analyses performed by the National Tidal Facility Australia of approximately eight years of Port Vila data, a rate of +10.5 mm per year has been observed (as compared to the global average, published by the IPCC, of around 1 or 2 mm per year). The Short Term Sea Level Trends plot shows how the trend estimate has varied over time, and because the data set is still relatively short, still varies considerably from month to month. In the early years, the trend appeared to indicate an enormous rate of sea level rise. Later, due to the 1997/1998 El Niño when sea level fell about 12 cm below average, the trend actually disappeared, and remained near zero for the next year. Over most of the past four years, the sea level trend appears to have remained close to its current value. We see that it is still far too early to deduce a long-term trend (or even whether it will be positive of negative) from this data.




June-2002

The sea level data recorded since installation is summarised in the following plot. The middle curve (green) represents the monthly mean sea level. The upper and lower curves show the highest and lowest values recorded each month. Unlike many of the SEAFRAME sites, sea level at Port Vila did not experience a dramatic decrease in 1998 as a result of El Niño, although (once seasonal effects are subtracted from the data) there was a fall of almost 25 cm between early 1997 and early 1998. Port Vila is relatively far from the equator, where El Niño signals are most pronounced.

By inspection of the monthly maxima (red curve) it appears that Vanuatu, like Fiji, the Cook Islands, Tonga and Tuvalu, experiences highest sea levels near the start of the year. At mid-year, the highest sea levels are typically about 20 cm less than when at the maximum. However, this pattern does not occur every year.


Monthly sea level at Port Vila SEAFRAME gauge



Historical Sea Level Trend Assessment

A longer sea level record is available at Vanuatu, from the Port Vila tide gauge operated from 1977 until 2000, with an eleven year gap - about 11 years in total of data. The University of Hawaii (UH) data exhibits a sea level rise of +6.21 mm/year over the period. The gauge was designed to monitor the variability caused by El Niño and shorter- term oceanic fluctuations, for which the high level of precision and datum control demanded by the determination of sea level trend were not required. Because of this, and the patchy data record, the trend can not be established without sizeable uncertainties.


Monthly sea level at Port Vila University of Hawaii data



Predicted highest astronomical tide

The component of sea level that is predictable due to the influence of the Sun and the Moon and some seasonal effects allow us to calculate the highest predictable level each year. It is primarily due to the ellipticity of the orbit of the Earth around the Sun, and that of the Moon around the Earth resulting in a point at which the Earth is closest to the Sun, combined with a spring tide in the usual 28 day orbit of the Moon around the Earth. The figure shows that the highest predicted level (1.58 m) over the period 1990 to 2016 will be reached at 17:28 Local Time on 24 November 2003.

Predicted highest tide each year for Port Villa




Monthly mean air temperature, water temperature and atmospheric pressure

The data summarised in the following three plots follows the same format as the preceding sea level plot: the middle curve (green) represents the monthly mean, and the upper and lower curves show the highest and lowest values recorded each month.


Compared to the more equatorial sites, Port Vila undergoes much greater seasonal temperature variations. The summertime highs are normally recorded in January or February. The minimum air temperature of 16.5°C was reached on 17 August 2001, and a maximum of 33.3°C was reached on 24 January 2002.

Monthly air temperature at Port Vila SEAFRAME gauge




Water temperature also undergoes seasonal oscillations, which are virtually in phase with those of air temperature. Interestingly, in several years the maxima in air and water temperature come a month or two after the sea level maxima.

Monthly water temperature at Port Vila SEAFRAME gauge



The sea level also responds to changes in barometric pressure. As a rule of thumb, a 1 hPa fall in the barometer, if sustained over a day or more, produces a 1 cm rise in the local sea level (within the area beneath the low pressure system). The seasonal (summertime) high sea levels at Port Vila are highly correlated with low barometric pressure systems. This is particularly the case for the very low pressure events (cyclones), most of which coincide with the highest sea levels for the year (since summer is also cyclone season). The lowest barometric pressures occurred on 29 March 1993 and 2 March 2001.

Monthly atmospheric pressure at Port Vila SEAFRAME gauge




June-2002

Extreme Events

Tropical Cyclones


A number of destructive Tropical Cyclones have passed near Vanuatu since the SEAFRAME was installed, and (as can be seen in the plot "Monthly atmospheric pressure at Lautoka") two in particular have come close enough to Port Vila to have been recorded as very low pressures. The first of these two was TC Prema, on 29 March 1993, and the second was TC Paula, on 2 March 2001. Both caused considerable damage. One consequence of TC Prema was that the SEAFRAME was damaged and inoperative for over a year and a half.



Cyclone map courtesy of Fiji Meteorological Service

June-2002

Track of Tropical Cyclone Paula, February/March 2001




Tsunami records


A tsunami can be defined as "A wave usually generated by seismic activity. Also called seismic sea wave, or, incorrectly, a *tidal wave*. Barely discernible in the open ocean, their amplitude may increase to over ten metres in the shallow coastal regions. Tsunamis are most common in the Pacific Ocean."

In June 2001 a tsunami was generated by an earthquake near Peru. The tsunami was detected at a number of SEAFRAME sites, with peak-to-trough amplitudes of over 20 cm at Lautoka, Fiji, for example. At Port Vila the amplitude was over 30 cm. Vanuatu is clearly not immune from potential problems should there be a large tsunami-generating undersea earthquake in the vicinity. The following plots show how, many hours after the initial earthquake, tsunamis can generate large disturbances in coastal locations.


Port Vila sea level in response to Peru Earthquake



Travel Times for Tsunami Wave from Peru Earthquake



Tsunami Wave due to Peru Earthquake (simulated magnitude)



June-2002


On 26 November 1999, 140 km to the northwest of Port Vila, a magnitude 7.3 (Mw) undersea earthquake occurred. The following is taken from the Center for International Disaster report::

The effects of the earthquake across Pentecost appear to be relatively localised. Eight fatalities have been reported; five persons died when their house collapsed in the central village of Ena; three others died after being swept out to sea by the tsunami. A further two are missing. About 40 serious injuries were also reported, particularly at Ena village, Melsisi Mission, Ranwadi High School, and the nearby areas. After the earthquake, the sea receded about 0.5 km from the shore. Some residents recognised this as the sign of an impending tsunami and managed to warn everyone to seek higher ground. The earthquake caused numerous landslides and, although destruction of garden areas is reported, crop losses are not thought to be widespread. In addition the tremor caused damage to both buildings and other infrastructure. The tsunami at Marteli Bay has flattened all low-lying structures including the brick schoolhouse, leaving only the church still standing. It destroyed the piped water supply, caused loss of all public and private possessions in the community including 7 boats and 4 generators, and covered gardens in silt. Banmatmat Bible College was also damaged, and landslides restrict the already limited access to the area.

We have added emphasis to part of the above paragraph, because it was reported that a number of lives were saved by the recognition of the warning signs for an impending tsunami.


The peak to trough amplitude at Port Vila was 0.9 m. The following graph is of the SEAFRAME data.

Port Vila sea level response to November 1999 Vanuatu Earthquake



At 17:22:49 on the 2nd of January 2002 (UTC) an earthquake magnitude 6.3mb (7.5MS) occurred at a depth of 21 km below the sea floor at 17.590S, 167.829E, 100 km west of Port Vila, Vanuatu. Several people were injured and there was widespread damage on the island of Efate. Access to the wharf was blocked by rockslides. The SEAFRAME tide gauge on the wharf at Port Vila recorded the tsunami wave that followed. Peak to trough amplitude reached 80 cm.


Port Vila sea level response to January 2002 Vanuatu Earthquake



<u>Definition of Datum and other Geodetic Levels at Port Vila, Vanuatu</u>

Newcomers to the study of sea level are confronted by bewildering references to "Chart Datum", "Tide Staff Zero", and other specialised terms. Frequent questions are, "how do NTFA sea levels relate to the depths on the marine chart?" and "how do the UH sea levels relate to NTFA's?".

Regular surveys to a set of coastal benchmarks are essential. If a SEAFRAME gauge or the wharf to which it fixed were to be damaged and need replacement, the survey history would enable the data record to be "spliced across" the gap, thereby preserving the entire invaluable record from start to finish.



The word "datum" in reference to tide gauges and nautical charts means a reference level. Similarly, when you measure the height of a child, your datum is the floor on which the child stands.

Where possible, "sea levels" in the NTFA data are normally reported relative to "Chart Datum" (CD), thus enabling users to relate the NTFA data (such as shown in the figure above) directly to depth soundings shown on marine charts – if the NTFA sea level is +1.5 metres, an additional 1.5 metres of water may be added to the chart sounding. At Port Vila, "LAT" (see below) provides an "equivalent" datum.

Mean Sea Level (MSL) in the figure is the average recorded level at the gauge over the year 1973. The MSL at Port Vila is 0.75 metres above CD.

Lowest Astronomical Tide, or "LAT", is based purely on tidal predictions over a 19 year period. In this case, LAT is 0.0 metres, meaning that if the sea level were controlled by tides alone, the sea level reported by NTFA would drop to 0.0 metres just once in 19 years.

